


This article was downloaded by: [Ms Kristin D. Neff]

On: 02 February 2015, At: 17:26

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



[Click for updates](#)

Self and Identity

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/psai20>

Self-Compassion, Stress, and Coping in the Context of Chronic Illness

Fuschia M. Sirois^{ab}, Danielle S. Molnar^c & Jameson K. Hirsch^d

^a Health and Well-Being Laboratory, Department of Psychology, Bishop's University, 2600 College St., Sherbrooke, Quebec, Canada

^b Centre de recherche sur le vieillissement, Sherbrooke, Quebec, Canada

^c Department of Psychology, Brock University, St. Catharines, Ontario, Canada

^d Laboratory of Rural Psychological and Physical Health, Department of Psychology, East Tennessee State University, Johnson City, TN, USA

Published online: 09 Jan 2015.

To cite this article: Fuschia M. Sirois, Danielle S. Molnar & Jameson K. Hirsch (2015): Self-Compassion, Stress, and Coping in the Context of Chronic Illness, *Self and Identity*, DOI: [10.1080/15298868.2014.996249](https://doi.org/10.1080/15298868.2014.996249)

To link to this article: <http://dx.doi.org/10.1080/15298868.2014.996249>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Self-Compassion, Stress, and Coping in the Context of Chronic Illness

Fuschia M. Sirois^{1,2}, Danielle S. Molnar³, and Jameson K. Hirsch⁴

¹Health and Well-Being Laboratory, Department of Psychology, Bishop's University, 2600 College St., Sherbrooke, Quebec, Canada

²Centre de recherche sur le vieillissement, Sherbrooke, Quebec, Canada

³Department of Psychology, Brock University, St. Catharines, Ontario, Canada

⁴Laboratory of Rural Psychological and Physical Health, Department of Psychology, East Tennessee State University, Johnson City, TN, USA

A recent review suggested that self-compassion promotes use of adaptive rather than maladaptive coping. Less is known about how self-compassion is linked to stress and coping in the context of a chronic stressor. Across two primarily female chronic illness samples, inflammatory bowel disease ($N = 155$) and arthritis ($N = 164$), a model linking self-compassion to lower stress through coping styles and coping efficacy was tested. Path analyses revealed significant indirect effects for adaptive coping styles (active, positive reframing, and acceptance), and negatively for maladaptive coping styles (behavioral disengagement and self-blame) in both samples. Findings suggest that the relative balance of adaptive and maladaptive coping strategies used by self-compassionate people is associated with better coping outcomes in the context of chronic illness.

Keywords: Self-compassion; Coping; Stress; Coping efficacy; Chronic illness.

Whether considered a momentary mindset or an enduring tendency, self-compassion is increasingly being recognized as an important quality for reducing stress. Self-compassion has been defined as a positive self-view that involves relating to oneself with kindness and acceptance in times of failure and difficulty (Neff, 2003b). A growing body of research has documented that self-compassion is linked to lower levels of perceived stress (Neely, Schallert, Mohammed, Roberts, & Chen, 2009; Sirois, 2014), and increased resilience in the face of stressful situations (Neff, Kirkpatrick, & Rude, 2007), including chronic illness (Brion, Leary, & Drabkin, 2014). In a recent theoretical review, Allen and Leary (2010) posited that self-compassionate people may experience lower stress because of their use of effective coping strategies, but found limited and mixed support for the hypothesis that people high in self-compassion prefer and use more adaptive, problem-focused coping styles and less maladaptive coping styles. Yet, there is little research on self-compassion, stress and coping in the context of an ongoing chronic stressor such as chronic disease.

Received 13 June 2014; accepted 3 December 2014; first published online 9 January 2015.

Gratitude is expressed to all the participants who took the time to complete the surveys. Preparation of this manuscript was supported by a Canada Research Chair awarded to the first author.

No potential conflict of interest was reported by the authors.

Correspondence should be addressed to: Fuschia M. Sirois, Department of Psychology, Bishop's University, 2600 College St., Sherbrooke, Quebec J1M 1Z7, Canada. Email: fsirois@ubishops.ca

Thus, it is unknown if the coping styles identified by Allen and Leary (2010) relate to self-compassion in this context. Understanding the potential of self-compassion for adaptive coping and therefore reducing stress in individuals with chronic disease, and especially chronic inflammatory disease, is an important goal in light of recent evidence highlighting the role of stress in the progression and exacerbation of inflammatory conditions (Cohen et al., 2012). In this study we aimed to address these issues and gaps in the research by testing the relations among self-compassion, adaptive and maladaptive coping styles, coping efficacy, and stress across two chronic illness groups.

Self-Compassion, Coping, and Stress

Self-compassion, as conceptualized by Neff (2003b), comprises three key features that may account for why self-compassionate people are able to successfully cope with stressful life circumstances. Self-kindness refers to a capacity to treat oneself with kindness and compassion rather than criticism or harsh self-judgment during challenging circumstances. Common humanity involves recognizing that painful and difficult experiences are part of the human condition rather than feeling isolated in one's suffering. Mindfulness involves taking a balanced perspective of one's emotions, particularly negative emotional states, rather than becoming over-identified and embroiled within these negative states (e.g., sadness, guilt, and anger). Together, these three qualities are proposed to reduce stress by helping individuals self-regulate the negative emotions that can arise from unexpected and/or uncontrollable events (Neff et al., 2007). In such instances, blaming events on one's own actions or failings can cyclically contribute to additional stress and impede healthy adjustment (Sirois, Davis, & Morgan, 2006). Thus, self-compassion may ameliorate stress by reducing coping that fosters negative emotional responses to stressors, and by promoting coping via adaptive behavioral or appraisal-based responses to stressors.

This view is consistent with appraisal-based models of coping and stress which emphasize the transactional nature of stress. Lazarus and Folkman (1984) cognitive transaction model of stress highlights the central role of the individuals' cognitive and behavioral responses in exacerbating or attenuating the stress response. Adaptive coping responses are those which successfully remove or reduce the stressor by cognitively changing its appraisal, or by making direct behavioral changes (Lazarus & Folkman, 1984). Thus, self-compassionate people may appraise stressors in a way that frames them as less negative and threatening, allowing them to engage in effective behavioral responses to reduce the stressor because their self-regulatory resources are not monopolized by the negative self-evaluations and mood that often arise in response to a stressful event (Sirois, Kitner, & Hirsch, *in press*; Terry & Leary, 2011).

Allen and Leary (2010) review of the evidence linking self-compassion to coping support this view. Using Skinner, Edge, Altman, and Sherwood (2003) taxonomy of coping as a conceptual lens they examined whether self-compassionate people were more likely to use adaptive coping strategies and less likely to use maladaptive coping strategies. Specific adaptive coping strategies reviewed were positive cognitive restructuring (e.g., reframing stressful events in a positive light), problem-solving (e.g., planning and taking active steps to deal with a stressor), support seeking (e.g., turning to others for support), distraction (e.g., reading or engaging in other distracting activities), whereas escape-avoidance coping strategies (e.g., cognitively or behaviorally disengaging from the stressor) were the less adaptive coping strategies reviewed. Overall, self-compassionate people were more likely to use positive cognitive restructuring and less likely to use escape-avoidance coping strategies. Evidence for links to problem-solving

was mixed. Some studies found links between self-compassion and active coping, planning and seeking instrumental support, and others did not; further, there was little evidence for links with seeking emotional support and distraction coping (Allen & Leary, 2010). They concluded that more research was needed to better understand how self-compassionate people handle stressors.

Self-Compassion and Coping with Chronic Illness

An important consideration missing from this preliminary theory and research on self-compassion and coping is whether these associations are relevant for dealing with chronic daily stressors, such as a chronic and functionally limiting illness. There is mounting evidence that ineffective management of stress in this context can negatively impact psychological and physical well-being (Evers et al., 2013; Maunder & Levenstein, 2008). This may be especially true for chronic inflammatory health conditions such as arthritis and inflammatory bowel disease (IBD) as stress can exacerbate inflammatory processes (Cohen et al., 2012).

The coping styles linked to self-compassion, as suggested by Allen and Leary (2010), may differentially impact the stress perceived by individuals with IBD and arthritis. Among patients with IBD, problem-focused and active coping strategies such as planning and instrumental support seeking are associated with better psychological outcomes (Graff et al., 2009; McCombie, Mulder, & Geary, 2013), whereas use of passive coping strategies are linked to worse outcomes (Jones, Wessinger, & Crowell, 2006). In patients with arthritis approach coping is linked to better psychological outcomes such as life satisfaction and lower psychological distress and depression (Treharne, Lyons, Booth, & Kitas, 2007; Vriezengkolk, van Lankveld, Geenen, & van den Ende, 2011). In contrast, escape-avoidance coping is associated with poor outcomes including psychiatric distress (Jones et al., 2006), helplessness and less illness acceptance (Voth & Sirois, 2009) in patients with IBD, and psychological distress in patients with arthritis (Ramjeet, Smith, & Adams, 2008; Vriezengkolk et al., 2011). Evidence is limited for positive reappraisals. Optimistic beliefs are linked to better adjustment outcomes in people with arthritis (Fournier, de Ridder, & Bensing, 2002). Similarly, acceptance, predicts better adjustment in both IBD (Voth & Sirois, 2009) and arthritis (Pinto-Gouveia, Costa, & Marôco, 2015). But, in the context of chronic illness distraction coping is not maladaptive as Allen and Leary (2010) suggest, but adaptive (Compas, Jaser, Dunn, & Rodriguez, 2012). For this reason it was not examined. Because evidence is mixed regarding whether emotion-focused coping strategies, including emotional support seeking, are associated with better or poorer adjustment in individuals with chronic illness (McCombie et al., 2013; Pellissier, Dantzer, Canini, Mathieu, & Bonaz, 2010; Schussler, 1992), we did not examine these strategies.

Self-blame is a coping strategy that may be very salient for understanding how self-compassion relates to adjustment in people with chronic illness. In the context of chronic illness, self-blame is associated with poor adjustment to chronic tinnitus (Sirois et al., 2006), and is linked to poor psychological well-being in patients with IBD (Voth & Sirois, 2009). Given that self-compassionate individuals treat themselves with kindness rather than harsh self-judgment in the face of a stressor, we expect that self-compassion will be negatively related to self-blame coping.

Individuals with chronic illness face a variety of daily and ongoing stressors, including pain and functional limitations, which can require using different coping strategies depending on the demand (Gignac, Cott, & Badley, 2000). Successful management of stress therefore relies less on the use of a single coping strategy, and more on the effectiveness of a set of coping strategies. Coping efficacy, appraisals of how successfully

one is coping with an illness-related stressor (Gignac et al., 2000), is one way to capture the degree to which a set of coping strategies are effective for managing stress in the context of chronic illness. For example, in a study of older adults with arthritis, higher coping efficacy was associated with less feelings of helplessness, dependence, and emotional reactivity (Gignac et al., 2000). In individuals with IBD, greater use of denial and behavioral disengagement coping is associated with lower coping efficacy (Voth & Sirois, 2009). Given the proposed links between self-compassion and the use of adaptive coping strategies outlined by Allen and Leary (2010), it is reasonable to expect that self-compassionate people with chronic illness would use a diverse set of coping strategies that would promote feeling that they are coping successfully with their illness, and in turn perceive less stress.

The Present Study

This study aimed to test and extend theory and research on self-compassion, coping and stress by examining their associations in individuals with IBD and arthritis. Consistent with previous research with healthy samples (Allen & Leary, 2010; Sirois, 2014), we hypothesized that self-compassion would relate to using effective coping strategies and lower perceived stress. We also expected that self-compassion would be positively linked to the adaptive coping strategies, and negatively linked to the maladaptive coping strategies noted by Allen and Leary (2010). We therefore examined a selected set of coping strategies theorized to be linked to self-compassion, and that research indicated a potential benefit for those with chronic illness: problem-solving coping (planning, instrumental support seeking, and active coping) and positive cognitive restructuring (positive reframing and acceptance), and maladaptive coping styles including escape avoidance coping (denial and behavioral disengagement) and self-blame.

Previous research has not directly examined the role of coping strategies with respect to stress among self-compassionate people. We therefore tested the efficacy of the coping strategies used by self-compassionate individuals and their associations with lower stress by examining the relative balance of adaptive and maladaptive coping strategies across the two illness groups using path analysis (see Figure 1). We posited that self-compassionate individuals in both samples would use a more adaptive set of coping strategies, which in turn would be associated with higher coping efficacy and, subsequently, would be associated with lower levels of stress.

Method

Participants and Procedure

The data analyzed for this study were collected as a 6-month follow-up to a study on self-perceptions and adjustment to illness among two chronic illness samples—individuals with any form of arthritis and individuals with IBD. Individuals who self-reported being medically diagnosed with either arthritis or IBD were eligible to participate; individuals




FIGURE 1 Conceptual model linking self-compassion to stress via coping styles and coping self-efficacy.

with an IBS, a related but distinct condition, were not eligible. Following clearance from the university research ethics board, participants were recruited via study notices placed in the community, on electronic support groups for people with arthritis and IBD, on the Arthritis Society research web page, and in the newsletter of the Crohn's and Colitis Foundation of Canada. The web page for each illness group directed participants to the corresponding online survey, housed on a secure university server. The Time 1 participation involved completing an online survey; there was no intervention involved. Participants consented to participate by clicking an "I agree" button on the online consent form, and were given the option to enter a drawing for a gift certificate to an online bookstore.

Total 325 people (170 with arthritis and 155 with IBD) completed the online survey at Time 2. Only 39.8% of the Time 1 participants with arthritis, and 36.2% of the Time 1 participants with IBD completed the Time 2 survey. Although participants were from a variety of locations around the world, the majority in both illness groups were from North America. **Table 1** presents the complete demographic characteristics of the two illness samples. Both samples were predominantly female and White. Among those with a self-reported diagnosis of any type of arthritis, rheumatoid arthritis (42.9%) and osteoarthritis (27.6%) were the most frequently reported subtypes, with fibromyalgia (7.1%), ankylosing spondylitis (5.9%), psoriatic arthritis (4.1%) and other subtypes (lupus, gout, and other arthritis types) also included. In the IBD group, 50.6% had Crohn's disease, 42.9% had ulcerative colitis, and 6.5% had unspecified colitis.

TABLE 1 Demographic Characteristics of Each Illness Sample

	Illness group	
	Arthritis	IBD
<i>N</i>	170	155
Sex (% female)	91.5	83.1
Age		
Mean (SD)	47.44 (11.6)	38.84 (12.8)
Range	18–75	18–72
Ethnicity (% Caucasian)	92.1	93.6
Country (%)		
Canada	49.1	48.7
United States	44.2	31.2
United Kingdom	4.12	14.3
Australia/New Zealand	1.2	1.3
Europe	1.2	4.5
Employment status (%)		
Full-time	36.4	43.6
Part-time	21	20.8
Unemployed/retired	18.5	25.5
Disabled	24.1	10.1
Education (%)		
High school or less	13.3	13.6
University or college	66.1	65.6
Graduate school	20.6	20.8
Relationship status (%)		
Married	57	67.5
Separated/divorced/widowed	20.6	7.9
Never married	22.4	24.5

Note: IBD, Inflammatory bowel disease; SD, standard deviation.

Measures

With the exception of disease-specific questions, participants completed identical surveys that included questions about demographic information, disease severity and duration, coping efficacy, coping, general perceived stress, and self-compassion. Scale means and reliabilities are presented in [Table 2](#). Although the coping measures were completed at Time 1 and 2, the self-compassion scale (SCS) was only completed at Time 2. Accordingly, the analyses focus on the Time 2 measures only.

Self-Compassion

Both samples completed the 26-item SCS (Neff, 2003a). The SCS assesses the three main components of self-compassion and their negative counterparts, self-kindness (self-judgment), common humanity (isolation), and mindfulness (over-identification). It includes both positively (“I try to be loving toward myself when I’m feeling emotional pain”) and negatively (“I’m disapproving and judgmental about my own flaws and inadequacies”) worded items reflecting the six components of self-compassion. Research indicates that the subscales are best explained by a single higher order factor of self-compassion as they are highly inter-correlated (Neff, 2003a). All items are prefaced with the statement “how I typically act towards myself during difficult times” and respondents indicate how often they behave in the described way using response options ranging from 1 (almost never) to 5 (almost always). Averaging the mean subscale scores after reverse coding the negative items yields a total self-compassion score. This scale has been successfully used in both student and community samples, demonstrating good validity, both convergent and discriminant, and excellent test-retest reliability previously ($\alpha = .93$) (Neff, 2003a; Neff & Pommier, 2013).

Disease-Related Variables

Participants indicated when they had been diagnosed with their IBD or arthritis. These dates were then transformed into years to create a time-since-diagnosis variable that was covaried in the main analyses. To assess the impact of their disease on daily living, participants rated the question “To what extent has arthritis affected your daily activities?” on a 4-point scale from 1 (not at all) to 4 (a lot). This item was chosen as a proxy for disease severity in both samples so that disease impact could be measured on a comparable scale.

Coping Strategies

The Brief COPE (Carver, 1997) is a well-validated, 30-item measure of the full COPE scale, that assesses how frequently functional and dysfunctional coping strategies are used. Each subscale is comprised of two items rated on a 4-point scale ranging from 1 (I usually do not do this at all) to 4 (I usually do this a lot). We examined eight subscales that previous theory and research suggest may be linked to both self-compassion and adjustment to chronic illness, including five adaptive coping strategies (instrumental support seeking, active coping, planning, positive reframing, and acceptance) and three maladaptive coping strategies (denial, behavioral disengagement, and self-blame) (Allen & Leary, 2010; Voth & Sirois, 2009). Participants reported their use of the coping strategies for dealing with their illness-related stress.

Coping-Efficacy

The three-item coping efficacy scale developed by Gignac and colleagues (Gignac et al., 2000) assessed appraisals of efficacy in coping with the chronic stressors associated with

TABLE 2 Descriptive Information and Correlations of the Model Variables in the Arthritis and IBD Samples

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13
1. SC	—	.19*	.41*	.33*	-.28*	.32*	.37*	-.35*	-.46*	.43*	-.56*	.10	.08
2. IN	.18*	—	.25*	.40*	.05	.28*	.21*	-.12	.15	.09	-.07	-.06	.14
3. ACT	.48*	.33*	—	.71*	-.16	.37*	.45*	-.34*	-.07	.40*	-.31*	.14	.01
4. PLAN	.41*	.35*	.63*	—	-.12	.51*	.60*	-.29*	.01	.36*	-.24*	.05	-.03
5. DEN	-.17*	.01	-.09	-.02	—	-.06	-.39*	.37*	.43*	-.35*	.31*	.15	.17*
6. PRF	.52*	.34*	.45*	.42*	.01	—	.36*	-.25*	.02	.35*	-.20*	.14	-.08
7. ACC	.28*	.12	.29*	.38*	-.26*	.29*	—	-.32*	-.21*	.48*	-.35*	.09	.05
8. BDIS	-.42*	.03	-.31*	-.16*	.32*	-.20*	-.17*	—	.37*	-.46*	.37*	-.09	.03
9. SB	-.50*	.17*	-.13	.03	.32*	.00	-.14	.51*	—	-.47*	.47*	.07	-.08
10. CSE	.47*	-.06	.38*	.23*	-.22*	.31*	.29*	-.56*	-.39*	—	-.54*	.01	.18*
11. PSS	-.56*	.02	-.35*	-.23*	.15	-.26*	-.10	.48*	.38*	-.57*	—	-.06	-.16
12. TDIAG	.16*	.04	.17*	.14	-.09	.10	.22*	-.24*	-.13	.25*	-.10	—	.24*
13. AGE	.29*	-.01	.06	.14	.06	.20*	.17*	-.13	-.15	.23*	-.26*	.25*	—
Mean (arthritis)	2.98	2.26	2.75	2.66	1.31	2.28	3.08	1.55	3.66	3.58	2.83	11.45	47.844
SD (arthritis)	.69	.81	.75	.83	.55	.87	.78	.74	1.70	.94	.84	10.36	11.56
Mean (IBD)	2.87	2.41	2.77	2.69	1.34	2.22	3.03	1.54	3.77	3.72	2.87	10.62	38.84
SD (IBD)	.71	.95	.88	.89	.56	.92	.80	.75	1.61	.91	.74	13.61	12.78
α (arthritis)	.94	.79	.60	.69	.44	.77	.75	.70	.71	.92	.90	—	—
α (IBD)	.94	.89	.81	.81	.53	.83	.80	.77	.67	.91	.91	—	—

Notes: Bivariate associations for the arthritis sample are presented in the lower diagonal and bivariate associations for the IBD sample are presented in the upper diagonal of the table. SC, self-compassion; IN, instrumental support coping; ACT, active coping; PLAN, planning coping; DEN, denial coping; PRF, positive reframing coping; ACC, acceptance coping; BDIS, behavioral disengagement coping; SB, self-blame coping; CSE, coping efficacy; PSS, perceived stress; TDIAG, time since diagnosis. * $p < .05$.

chronic illness. The scale focuses on three common challenges associated with chronic illness: symptoms, emotional aspects, and day-to-day problems. Items such as “I am successfully coping with the symptoms of my arthritis” are scored on a 5-point Likert type scale with responses ranging from 1 (strongly disagree) to 5 (strongly agree); higher scores reflect greater coping efficacy. For the IBD sample, the word “arthritis” in all items was replaced with “IBD.”

Perceived Stress

Each sample completed the 10-item version of the Perceived Stress Scale (PSS; Cohen & Williamson, 1988), a widely used, empirically established index of general stress. The perceived stressfulness of events experienced within the past month is assessed with items such as “In the last month, how often have you felt nervous and stressed” rated on a 5-point scale with response options ranging from “never” to very “often.” The PSS has demonstrated good internal consistency in previous studies (Cohen & Williamson, 1988), and in the current samples (see Table 2).

Statistical Analyses

Differences between the two samples were evaluated using *t*-tests. With correlation analyses, we tested the proposed links between self-compassion, coping strategies, self-efficacy and stress. In multivariate analyses, we also tested the hypothesis that self-compassion was associated with a relative balance of more adaptive coping strategies and less maladaptive coping strategies and, in turn, to better coping efficacy and lower stress across both chronic illness samples (Figure 1), using path analysis via Mplus 7.11 (Muthen & Muthen, 2013). First, we examined the overall model combining both the arthritis and IBD samples (see Table 2 top panel for summary). Full-information maximum likelihood (Arbuckle, 1996) estimation procedures were employed and the standardized estimates are presented. Model fit was assessed using the comparative fit index (CFI), the root-mean-square error of approximation (RMSEA), and the standardized root mean square residual (SRMR). Indirect effects were tested using the biased-corrected bootstrap method with 1000 resamples and the 95% bias-corrected confidence intervals (CIs). This method provides a more accurate balance between Type 1 and Type 2 errors compared with other methods used to test indirect effects (MacKinnon, Lockwood, & Williams, 2004). Finally, the consistency of the path model results across sample (arthritis versus IBD) was assessed using multiple group analysis: model fit indices were compared between a path model in which all predictive paths were freely estimated within the two samples, and a constrained model, in which corresponding predictive paths were set equal across samples. Small, non-significant changes in the model fit indices indicate that the assumption of equality in corresponding predictive paths across samples was tenable (Kline, 2005).

Results

Descriptive Information

Descriptive information along with correlations between all study variables, for each sample, is presented in Table 2. All variables were normally distributed except for denial (skewness = 1.96, SE = .14), and behavioural disengagement coping (skewness = 1.47, SE = .14), which were positively skewed. Across both samples, self-compassion was positively associated with all adaptive coping styles and coping efficacy, and negatively associated with the maladaptive coping styles and perceived stress. Results from

independent sample *t*-tests revealed that there were no sample-related differences in scores on model variables, except with respect to age $t(315) = 6.41, p < .001$; individuals with arthritis were significantly older than those with IBD (see [Table 1](#)). Consequently, age was included in the analyses as a covariate. Given the heterogeneity of symptoms across the two samples, we used effects coding and included a set of seven variables in the analyses to account for differences in symptoms across samples. This allowed us to determine the effect of being in a particular subgroup (i.e., rheumatoid arthritis, osteoarthritis, fibromyalgia, ankylosing spondylitis, other arthritic conditions, Crohn's disease, and ulcerative colitis) relative to the remaining sample with respect to the model variables.

Path Analyses

The conceptual model depicted in [Figure 1](#), accounting for age and heterogeneity of symptoms across samples, was tested. Goodness of fit indices revealed that the structure of the hypothesized model did not provide an acceptable explanation of the data ($\chi^2_{(10)} = 68.35, p < .001, CFI = .95, RMSEA = .14, p_{close} < .001, SRMR = .03$). Modification indices indicated that adding a direct path from self-compassion to stress would significantly improve model fit. The addition of this path greatly improved model fit ($\chi^2_{diff(1)} = 56.05, p < .001, \Delta CFI = .05, \Delta RMSEA = .11$) and resulted in a model that fit the data well ($\chi^2_{(9)} = 12.30, p = .20, CFI = 1.00, RMSEA = .03, p_{close} = .68, SRMR = .01$). The results from the final model, accounting for 43% of the variance in stress, are presented in [Figure 2](#). As expected, coping self-efficacy was negatively associated with stress. Active coping, positive reframing, and acceptance were positively associated with coping self-efficacy, whereas behavioral disengagement and self-blame were negatively associated with coping self-efficacy. Self-compassion was positively associated with instrumental coping, active coping, planning, positive reframing, and acceptance, and negatively associated with denial, behavioral disengagement, and self-blame. The illness subtype comparisons revealed that fibromyalgia was associated with less coping self-efficacy, whereas ulcerative colitis and Crohn's disease were associated with higher levels of coping self-efficacy relative to others. Individuals with ankylosing spondylitis used less planning relative to others. Those with fibromyalgia were more likely to use positive re-framing, and those with ulcerative colitis were less likely to use behavioral disengagement compared with the rest of the sample.

In addition to the direct negative association between self-compassion and stress, self-compassion also shared significant and negative indirect associations with stress via active coping and coping self-efficacy [$b = -.030; 95\% CI = (-.060; -.005)$], positive reframing and coping self-efficacy [$b = -.033; 95\% CI = (-.062; -.016)$], acceptance and coping self-efficacy [$b = -.020; 95\% CI = (-.041; -.006)$], behavioral disengagement and coping self-efficacy [$b = -.038; 95\% CI = (-.077; -.018)$], and through self-blame and coping self-efficacy [$b = -.048; 95\% CI = (-.084; -.025)$]. Concerning the consistency of results across samples, the assumption of equality across illness groups was tenable ($\Delta\chi^2 = 55.11, df = 56, p = .50; \Delta CFI = .001; \Delta RMSEA = .019$).

Discussion

Across two chronic illness samples we found that self-compassion was associated with greater use of adaptive coping and less use of maladaptive coping which, in turn, were linked to coping efficacy and subsequently less stress. After accounting for indirect relationships, the direct relationship between self-compassion and stress remained significant, indicating that self-compassion is linked to lower stress through routes beyond




FIGURE 2 Final model linking self-compassion to stress via coping styles and coping self-efficacy. *Notes:* Statistically significant paths are shown via solid lines, whereas the dashed lines represent paths that were estimated in the model, but were not statistically significant. A set of seven variables that were created using effects coding were included in the analyses to account for differences in symptoms among the different samples, but not shown for ease of presentation. Respondent's age was also accounted for in the analyses, but not shown for ease of presentation. Covariances between disturbances associated with coping styles were estimated, but not shown for ease of presentation. Standardized path coefficients are displayed. * $p < .05$.

copied. The model did not differ across the two illness samples, suggesting that self-compassion is a quality associated with adjustment to stressors in the context of IBD and arthritis.

Our findings replicate and extend previous research on self-compassion and coping in several important ways. Whereas past research has demonstrated mixed findings with respect to self-compassion and problem-solving coping styles (Allen & Leary, 2010), we found associations with three coping styles in this category at the bivariate level: instrumental support seeking, active coping and planning, all of which are cognitive or behavioral in nature. However, only active coping linked self-compassion to coping efficacy, indicating that this problem-focused coping strategy may be more effective for coping with IBD and arthritis. Findings regarding the use of positive cognitive restructuring strategies (positive reframing and acceptance) were consistent with previous research (Allen & Leary, 2010); however, we found that these strategies linked self-compassion to better coping efficacy and, in turn, to less stress. In the context of chronic inflammatory illness, research suggests that acceptance—a cognitive appraisal process—may be especially important for managing day-to-day stress and promoting adjustment (Pinto-Gouveia et al., 2015; Ramjeet, Koutantji, Barrett, & Scott, 2005; Voth & Sirois, 2009). Thus, self-compassionate people may have a coping advantage that promotes adjustment to illness because they reframe how they view illness-related challenges and, therefore, engage in appropriate, rather than unrealistic, problem-solving strategies to minimize stress.

As expected, self-compassion was negatively related to each of the escape-avoidance coping strategies, denial and behavioral disengagement. Previous research has noted a negative association between self-compassion and trait procrastination, a form of behavioral disengagement and avoidant coping (Sirois, 2014). Consistent with this, we found that less use of behavioral disengagement was linked to successful coping and less stress. The negative association with self-blame coping is unique to this study and it highlights the importance of the self-kindness dimension of self-compassion (Neff, 2003b) for minimizing harsh and self-critical thoughts when dealing with illness-related stressors. In low control circumstances such as chronic illness, it is not uncommon for individuals to try to regain control by retrospectively attributing uncontrollable illness-related changes to their own behavior (Thompson, Sobolew-Shubin, Galbraith, Schwankovsky, & Cruzen, 1993) and view these changes as personal failures. This self-blaming response is linked to poor adjustment in other chronic conditions (Sirois et al., 2006; Thompson, Cheek, & Graham, 1988), and poor coping efficacy in IBD (Voth & Sirois, 2009). Our findings provide preliminary and suggestive evidence that self-compassion may be a protective factor against this form of maladaptive coping.

The current findings also demonstrate, for the first time, that it is the relative balance of the greater use of adaptive and less use of maladaptive coping strategies that links self-compassion to better coping outcomes. In their review of current research, Allen and Leary (2010) posited that this may be the case. By using a path analysis model that allowed for simultaneous testing of the associations of self-compassion with several adaptive and maladaptive coping strategies, we were able to provide more direct empirical support for this proposition and were also able to examine how the coping styles were linked to coping efficacy and lower stress in our health-compromised samples. Although cross-sectional, our analyses suggest that self-compassionate individuals with chronic illness may have a repertoire of adaptive coping styles that they call upon to deal with different stressors associated with their illness. This view of self-compassionate individuals is consistent with theory and research on coping flexibility which highlights the benefits of being able to shift among different coping strategies to find the most effective response for psychological health and well-being (Kato, 2012). Examining how self-compassion relates to flexible coping may, therefore, be a fruitful area for future research.

When considered in light of the predominantly female sample, the links between self-compassion and coping in this study both parallel and contrast research on how women cope with stress in comparison with men. For example, a meta-analysis of sex differences in coping found that women in comparison with men tend to use positive self-talk to cope with stressors (Tamres, Janicki, & Helgeson, 2002). As noted previously, this positive reappraisal style of coping reflects the tendency of self-compassionate people to reframe stressors as less threatening. However, this meta-analysis and other research found that women tend to use more emotion-focused strategies, including rumination, as well as more avoidant coping strategies, than men (Matud, 2004; Tamres et al., 2002). In this study self-compassionate individuals with IBD and arthritis tended to use less avoidant coping, and less self-blame coping, a strategy akin to rumination in that it involves repetitive, unproductive thinking that is associated with greater feelings of distress and less self-compassion (Filip, 2010). Taken together, the current findings and research highlighting how women cope with stress provides suggestive evidence that self-compassion may be especially beneficial for women coping with the stressors of chronic illness because it promotes the use of adaptive coping strategies while minimizing the use of maladaptive coping strategies.

Despite being novel, the results from this study should be considered in light of several limitations. Although, the proposed links among self-compassion, coping, coping efficacy and stress are informed by theory and experimental research (Allen & Leary, 2010; Neff, 2003b), the cross-sectional nature of our data precludes any strong conclusions about causality. To determine the temporal precedence of the model it should be tested within the framework of a prospective longitudinal study. We tested a limited number of coping styles as suggested by theory and research; however, future research could provide a more comprehensive portrait of the repertoire of coping styles used by self-compassionate individuals in the context of chronic illness. It should also be noted that the direct association from self-compassion to stress remained significant, highlighting the role of factors, coping-related or otherwise, not tested in this study that might explain why self-compassionate people perceive less stress. For example, the mindfulness component of self-compassion, with its emphasis on not becoming embroiled in negative mood states, may account for this link. Further research on understanding the direct link between self-compassion and stress would bring clarity to this issue.

Although our findings are generally consistent with previous research in both healthy and health-compromised individuals, the coping styles associated with self-compassion in this study may not apply to other samples. Indeed, the diversity of sub-types within each illness group makes it difficult to determine if the results apply more or less to any one subtype. Moreover, IBD and arthritis, though both forms of chronic inflammatory disease, are also very distinct in many ways, despite the results suggesting that the relations among the study variables did not differ significantly between the two groups, or as a function of any one subtype. Both samples were primarily White and female also limiting the applicability of the findings to more diverse samples. The results may also be specific to chronic inflammatory conditions and may not generalize to other chronic illness groups or to general medical populations. Collectively, these limitations indicate that more work with other illness populations would clarify the extent to which the findings can be generalized to other samples. Nonetheless, this study is the first that we are aware of to examine self-compassion and coping with IBD and arthritis.

This study replicated and extended previous theory and research, finding that in women with IBD and arthritis, self-compassion is linked to greater use of adaptive coping and less use of maladaptive coping for illness-related stressors, which, in turn, was linked to successful coping and less stress. The healthy repertoire of coping styles associated with self-compassion found in this study provides suggestive evidence that cultivating a tendency to relate to oneself with kindness, compassion and acceptance in times of failure and difficulty may be valuable for reducing stress for those coping with chronic inflammatory disease.

References

- Allen, A. B., & Leary, M. R. (2010). Self-compassion, stress, and coping. *Social and Personality Psychology Compass*, 4, 107–118.
- Arbuckle, J. L. (1996). Full information estimation in the presence of incomplete data. In G. A. Marcoulides & R. E. Schumacker (Eds.), *Advanced structural equation modeling: Issues and techniques* (Vol. 243–277). Mahwah, NJ: Lawrence Erlbaum Associates.
- Brion, J., Leary, M., & Drabkin, A. (2014). Self-compassion and reactions to serious illness: The case of HIV. *Journal of Health Psychology*, 19, 218–229.
- Carver, C. S. (1997). You want to measure coping but your protocol's too long: Consider the brief COPE. *International Journal of Behavioral Medicine*, 4, 92–100.

- Cohen, S., Janicki-Deverts, D., Doyle, W. J., Miller, G. E., Frank, E., Rabin, B. S., & Turner, R. B. (2012). Chronic stress, glucocorticoid receptor resistance, inflammation, and disease risk. *Proceedings of the National Academy of Sciences*, *109*, 5995–5999.
- Cohen, S., & Williamson, G. (1988). Perceived stress in a probability sample of the United States. In S. Spacapan & S. Oskamp (Eds.), *The social psychology of health: Claremont symposium on applied social psychology*. Newbury Park, CA: Sage.
- Compas, B. E., Jaser, S. S., Dunn, M. J., & Rodriguez, E. M. (2012). Coping with chronic illness in childhood and adolescence. *Annual Review of Clinical Psychology*, *8*, 455–480.
- Evers, A. W. M., Verhoeven, E. W. M., van Middendorp, H., Sweep, F. C. G. J., Kraaijmaat, F. W., Donders, A. R. T., ... van Riel, P. L. C. M. (2013). Does stress affect the joints? Daily stressors, stress vulnerability, immune and HPA axis activity, and short-term disease and symptom fluctuations in rheumatoid arthritis. *Annals of the Rheumatic Diseases*, *73*, 1683–1688.
- Filip, R. (2010). Rumination and worry as mediators of the relationship between self-compassion and depression and anxiety. *Personality and Individual Differences*, *48*, 757–761.
- Fournier, M., de Ridder, D., & Bensing, J. (2002). Optimism and adaptation to chronic disease: The role of optimism in relation to self-care options of type 1 diabetes mellitus, rheumatoid arthritis and multiple sclerosis. *British Journal of Health Psychology*, *7*, 409–432.
- Gignac, M. A. M., Cott, C., & Badley, E. M. (2000). Adaptation to chronic illness and disability and its relationship to perceptions of independence and dependence. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, *55*, 362–372.
- Graff, L. A., Walker, J. R., Clara, I., Lix, L., Miller, N., Rogala, L., ... Bernstein, C. N. (2009). Stress coping, distress, and health perceptions in inflammatory bowel disease and community controls. *American Journal of Gastroenterology*, *104*, 2959–2969.
- Jones, M. P., Wessinger, S., & Crowell, M. D. (2006). Coping strategies and interpersonal support in patients with irritable bowel syndrome and inflammatory bowel disease. *Clinical Gastroenterology and Hepatology*, *4*, 474–481.
- Kato, T. (2012). Development of the Coping Flexibility Scale: Evidence for the coping flexibility hypothesis. *Journal of Counseling Psychology*, *59*, 262–273.
- Kline, R. B. (2005). *Principles and practice of structural equation modeling* (2nd ed.). New York, NY: The Guilford Press.
- Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping*. New York, NY: Springer.
- MacKinnon, D. P., Lockwood, C. M., & Williams, J. (2004). Confidence limits for the indirect effect: Distribution of the product and resampling methods. *Multivariate Behavioral Research*, *39*, 99–128.
- Matud, M. P. (2004). Gender differences in stress and coping styles. *Personality and Individual Differences*, *37*, 1401–1415.
- Maunder, R. G., & Levenstein, S. (2008). The role of stress in the development and clinical course of inflammatory bowel disease: Epidemiological evidence. *Current Molecular Medicine*, *8*, 247–252.
- McCombie, A. M., Mulder, R. T., & Geary, R. B. (2013). How IBD patients cope with IBD: A systematic review. *Journal of Crohn's and Colitis*, *7*, 89–106.
- Muthen, L. K., & Muthen, B. O. (2013). *Mplus user's guide*. Los Angeles, CA: Muthen and Muthen.
- Neely, M., Schallert, D., Mohammed, S., Roberts, R., & Chen, Y.-J. (2009). Self-kindness when facing stress: The role of self-compassion, goal regulation, and support in college students' well-being. *Motivation and Emotion*, *33*, 88–97.
- Neff, K. D. (2003a). Development and validation of a scale to measure self-compassion. *Self and Identity*, *2*, 223–250.
- Neff, K. D. (2003b). Self-compassion: An alternative conceptualization of a healthy attitude toward oneself. *Self and Identity*, *2*, 85–101.
- Neff, K. D., Kirkpatrick, K. L., & Rude, S. S. (2007). Self-compassion and adaptive psychological functioning. *Journal of Research in Personality*, *41*, 139–154.

- Neff, K. D., & Pommier, E. (2013). The relationship between self-compassion and other-focused concern among college undergraduates, community adults, and practicing meditators. *Self and Identity, 12*, 160–176.
- Pellissier, S., Dantzer, C., Canini, F., Mathieu, N., & Bonaz, B. (2010). Psychological adjustment and autonomic disturbances in inflammatory bowel diseases and irritable bowel syndrome. *Psychoneuroendocrinology, 35*, 653–662.
- Pinto-Gouveia, J., Costa, J., & Marôco, J. (2015). The first 2-years of rheumatoid arthritis: The influence of acceptance on pain, physical limitation and depression. *Journal of Health Psychology, 20*, 102–112.
- Ramjeet, J., Koutantji, M., Barrett, E. M., & Scott, D. G. I. (2005). Coping and psychological adjustment in recent-onset inflammatory polyarthritis: The role of gender and age. *Rheumatology, 44*, 1166–1168.
- Ramjeet, J., Smith, J., & Adams, M. (2008). The relationship between coping and psychological and physical adjustment in rheumatoid arthritis: A literature review. *Journal of Clinical Nursing, 17*, 418–428.
- Schussler, G. (1992). Coping strategies and individual meanings of illness. *Social Science and Medicine, 34*, 427–432.
- Sirois, F. M. (2014). Procrastination and stress: Exploring the role of self-compassion. *Self and Identity, 13*, 128–145.
- Sirois, F. M., Davis, C. G., & Morgan, M. S. (2006). “Learning to live with what you can’t rise above”: Control beliefs, symptom control, and adjustment to tinnitus. *Health Psychology, 25*, 119–123.
- Sirois, F. M., Kitner, R., & Hirsch, J. K. (in press). Self-compassion, affect, and health behaviors. *Health Psychology*.
- Skinner, E. A., Edge, K., Altman, J., & Sherwood, H. (2003). Searching for the structure of coping: A review and critique of category systems for classifying ways of coping. *Psychological Bulletin, 129*, 216–269.
- Tamres, L. K., Janicki, D., & Helgeson, V. S. (2002). Sex differences in coping behavior: A meta-analytic review and an examination of relative coping. *Personality and Social Psychology Review, 6*, 2–30.
- Terry, M. L., & Leary, M. R. (2011). Self-compassion, self-regulation, and health. *Self and Identity, 10*, 352–362.
- Thompson, S. C., Cheek, P. R., & Graham, M. A. (1988). The other side of perceived control: Disadvantages and negative effects. In S. Spacapan & S. Oskamp (Eds.), *The Social Psychology of Health* (pp. 69–93). Newbury Park, CA: Sage.
- Thompson, S. C., Sobolew-Shubin, A., Galbraith, M. E., Schwankovsky, L., & Cruzen, D. (1993). Maintaining perceptions of control: Finding perceived control in low-control circumstances. *Journal of Personality and Social Psychology, 64*, 293–304.
- Treharne, G. J., Lyons, A. C., Booth, D. A., & Kitas, G. D. (2007). Psychological well-being across 1 year with rheumatoid arthritis: Coping resources as buffers of perceived stress. *British Journal of Health Psychology, 12*, 323–345.
- Voth, J., & Sirois, F. M. (2009). The role of self-blame and responsibility in adjustment to inflammatory bowel disease. *Rehabilitation Psychology, 54*, 99–108.
- Vriezekolk, J. E., van Lankveld, W. G., Geenen, R., & van den Ende, C. H. (2011). Longitudinal association between coping and psychological distress in rheumatoid arthritis: A systematic review. *Annals of the Rheumatic Diseases, 70*, 1243–1250.